

ASPB Pioneer Member

Mary Helen Goldsmith

Mary Helen Goldsmith (née Martin), curious about and loving nature, grew up in Maryland. She earned a B.A. degree in Zoology and Chemistry from Cornell University in 1955, and a Ph.D. from Radcliff College at Harvard University in 1959. She studied at Harvard with Professor Kenneth V. Thimann, one of the world's authorities on auxin, at a time when how auxin moved through tissues was a greatly motivating mystery in plant physiology. How could a molecule related to tryptophan move so readily through tissue in one direction but barely budge in the other direction?¹ Why was its transport through tissue, but not its initial uptake into cells, so reliant on respiration?² Mary Helen continued to investigate these puzzles at Yale University as her husband began his neuroscience career there. Mary Helen's position at Yale was first a Research Associate (1961–1973), then a Lecturer (1963–1973), and finally (1984) a tenured Professor.

In the 1960s, when corn and oat coleoptiles were the model systems, Mary Helen's research relied on ¹⁴C-labeled IAA applied as a pulse to track auxin through tissues³. In the 1970s, Mary Helen's research began to feature micro-electrode measurements of electric potential differences across membranes, because knowledge of these and other electrophysiological parameters was needed



to better understand auxin transport^{4,5}. Mary Helen synthesized the field's and her results in an influential review article⁶ that Bob Cleland (University of Washington) said was the paper that "brought the chemiosmotic hypothesis of PAT [polar auxin transport] to the attention of auxin researchers" (Figure 1). In this paper, Mary Helen clearly explained how protons pumps, pH gradients, the membrane potential, and polarly localized anionic auxin-permeable channels (not yet discovered) could form a mechanism that moves the hormone through plants from cell to cell. Mary Helen's careful measurements of auxin transport velocities provided the data needed to parameterize a model that produced an important result: in order for the entire tissue to display the measured degree of bias and speed of auxin transport, the downstream end of each cell in a file needs to be only about two-fold more permeable to the

auxin anion than the opposite end. This paper⁷, authored by Mary Helen, her husband, and her father (a renowned mathematician at the University of Maryland), should be read today by all researchers studying PIN and ABCB auxin transporters.

In the 1980s, more so than measuring polar transport through tissues, Mary Helen used radio-tracer and electrophysiological methods to study auxin-related membrane physiology. She incorporated biochemical methods to investigate auxin uptake and the binding of pharmacological agents that impair transport. Using plasma membrane vesicles and coleoptile sections^{8,10}, she obtained electrophysiological support for the idea that auxin stimulated the plasma membrane proton pump^{11,12}, a key tenet of the acid growth hypothesis. When, in the 1990s, the patch clamp technique promised to revolutionize the study of plant membrane transport, Mary Helen brought this new technology into her lab¹³. At the same time, Arabidopsis was replacing traditional systems for studying almost everything. Using the patch clamp technique¹⁴, Mary Helen embraced both developments and became among the first to characterize ion channels in Arabidopsis plasma membranes.

Throughout her career, Mary Helen served the American Society of Plant Physiologists as an elected member of the Executive Committee, a member of the Board

continued on next page

ASPB Pioneer Member

Mary Helen Goldsmith *continued*



of Trustees, a *Plant Physiology* editor, and a member of various committees. In 1990, Mary Helen was elected President of the society, only the second woman to hold the office.

The rigorous thinking and hard work needed to generate new knowledge in auxin transport physiology never diminished Mary Helen's love of nature. Instead, she nourished it in her job as the director of Yale's Marsh Botanical Garden, and she made it possible for countless Yale students to develop their own genuine connection with the living world by establishing the Environmental Studies program at Yale, which is now a complete undergraduate major. As of April 2022, Mary Helen is living with her husband Tim in Hamden, Connecticut, with plenty of reasons to feel satisfied.

References

- Goldsmith, M.H.M. and Thimann, K.V.** (1962). Some characteristics of movement of indoleacetic acid in coleoptiles of *Avena*. I. Uptake, destruction, immobilization, & distribution of IAA during basipetal translocation. *Plant Physiology* **37**: 492-505
- Goldsmith, M.H.M.** (1967). Separation of transit of auxin from uptake: average velocity and reversible inhibition by anaerobic conditions. *Science* **156**: 661-663
- Goldsmith, M.H.M.** (1967). Movement of pulses of labeled auxin in corn coleoptiles. *Plant Physiology* **42**: 258-263
- Goldsmith, M.H.M., Fernández, H.R. and Goldsmith, T.H.** (1972). Electrical properties of parenchymal cell membranes in the oat coleoptile. *Planta* **102**: 302-323

Bob Cleland (University of Washington) on Mary Helen Goldsmith's contributions



Mary Helen Goldsmith at her retirement symposium, Yale University, 2006

Figure 1. Bob Cleland (University of Washington) spoke about Mary Helen's career at her retirement symposium at Yale University in 2006.

5. Goldsmith, T.H. and Goldsmith, M.H.M. (1978). The interpretation of intracellular measurements of membrane potential, resistance, and coupling in cells of higher plants. *Planta* **143**: 267-274

6. Goldsmith, M.H.M. (1977). The polar transport of auxin. *Annual Review of Plant Physiology* **28**: 439-478

7. Goldsmith, M.H.M., Goldsmith, T.H. and Martin, M.H. (1981) Mathematical analysis of the chemosmotic polar diffusion of auxin through plant tissues. *Proceedings of the National Academy of Sciences USA*. **78**: 976-980

8. Edwards, K.L. and Goldsmith, M.H.M. (1980). pH-Dependent accumulation of indoleacetic acid by corn coleoptile sections. *Planta* **147**: 457-466

9. Sussman, M.R. and Goldsmith, M.H.M. (1981). The action of specific inhibitors of auxin transport on uptake of auxin and binding of N-1-naphthylphthalamic acid to a membrane site in maize coleoptiles. *Planta* **152**: 13-18

10. Clark, K.A. and Goldsmith, M.H.M. (1987). Effect of surface and membrane potentials on IAA uptake and binding by Zucchini membrane vesicles. In:

Klämbt, D. (eds) *Plant Hormone Receptors*. NATO ASI Series, vol 10. Springer, Berlin, Heidelberg.

11. Bates, G.W. and Goldsmith, M.H.M. (1983). Rapid response of the plasma-membrane potential in oat coleoptiles to auxin and other weak acids. *Planta* **159**: 231-237

12. Senn, A.P. and Goldsmith, M.H.M. (1988) Regulation of electrogenic proton pumping by auxin and fusicoccin as related to the growth of *Avena* coleoptiles. *Plant Physiology* **88**: 131-138

13. Kourie J. and Goldsmith, M.H.M. (1992). K⁺ channels are responsible for an inwardly rectifying current in the plasma membrane of mesophyll protoplasts of *Avena sativa*. *Plant Physiology* **98**: 1087-1097

14. Spalding, E.P. and Goldsmith, M.H.M. (1993). Activation of K⁺ channels in the plasma membrane of *Arabidopsis* by ATP produced photosynthetically. *The Plant Cell* **5**: 477-484